


EMBRYOLOGIA

VOLUME 5


1960-1961

THE EMBRYOLOGIA SOCIETY

Nagoya, Japan



M39845



RECEIVED
JAN 10 1964
U.S. AIR FORCE
HONOLULU, HAWAII

EMBRYOLOGIA

TABLE OF CONTENTS

VOLUME 5

No. 1 (April, 1960)

	PAGE
KOJIMA, M. K.: Cyclic changes of the cortex and the cytoplasm of the fertilized and the activated sea urchin egg.	
I. Changes in the thickness of the hyaline layer.....	1
TAKATA, C. and YAMADA, T.: Endodermal tissues developed from the isolated newt ectoderm under the influence of guinea pig, bone marrow.	8
DAN, K. and KUBOTA, H.: Data on the spawning of <i>Comanthus japonica</i> between 1937 and 1955.	21
TAKATA, C.: The differentiation <i>in vitro</i> of the isolated endoderm under the influence of the mesoderm in <i>Triturus pyrrhogaster</i>	38
KOJIMA, M. K.: The effect of DNP and NaN_3 on fertilized eggs of the sea urchin with special reference to the induction of the abnormal cleavage.	71
TAKAOKA, M.: Studies of the metamorphosis in insects. IV. Inhibition of pupation by carbon dioxide in the mature larva of <i>Drosophila melanogaster</i>	78
ISHIKAWA, M.: The effects of sulfhydryl reagents on artificial parthenogenesis in the sea urchin egg.....	85
MURAMATSU, S.: Studies on the physiology of <i>Artemia</i> embryos. I. Respiration and its main substrate during the early development of the encysted embryo.	95
KAJISHIMA, T.: Analysis of gene action in the transparent-scaled goldfish, <i>Carassius auratus</i> . I. On the gene action in the disappearance of guanophores.	107

No. 2 (July, 1960)

- KAJISHIMA, T.: Analysis of gene action in the transparent-scaled goldfish, *Carassius auratus*. II. The effects of pituitary and thyroid on gene action.127
- PFAUTSCH, M.-E.: Untersuchung des Nukleinsäuregehaltes in verschiedenen Keimregionen bei der frühen Gastrula und Neurula von *Triturus alpestris* (LAUR) und *Ambystoma mexicanum* (COPE).139
- TAKEUCHI, K.: The behavior of carotenoid and distribution of xanthophores during development of the medaka (*Oryzias latipes*).170
- KOJIMA, M. K.: Cyclic changes of the cortex and the cytoplasm of the fertilized and the activated sea urchin egg. II. The formation of clear spots by the hypertonicity, ether and urethane.178
- ISHIKAWA, M.: Experimental formation of the mark of sperm-entry on the fertilization membrane in the sea urchin egg.186
- TAKATA, C.: The differentiation *in vitro* of the isolated endoderm in the presence of the neural fold in *Triturus pyrrhogaster*.194
- EGUCHI, Y.: Experimental studies on the adrenal cortex of the mouse-fetus. I. Effects of maternal adrenalectomy on the adrenal of the fetus based on histology and volume determination.206

No. 3 (December, 1960)

- EGUCHI, G. and ISHIKAWA, M.: ³²P-Uptake by dorsal and ventral halves of the iris at the beginning of lens regeneration in the newt.219
- TOMITA, H.: On the nature of amphibian phenol oxidase.227
- ANDO, S.: Physiological study on egg formation of the fish. I. Accumulation of carbohydrates and proteins during oogenesis.239
- SAWADA, N.: Studies on the artificial activation and cortical reaction of the egg of *Nereis japonica*.247
- TAKAOKA, M.: Studies of the metamorphosis in insect. V.

Factors controlling the larval period of the squash fly, <i>Zeugoducus depressus</i> SHIRAKI.....	259
ŌISHI, S.: Studies on the teloblasts in the decapod embryo. II. Origin of teloblasts in <i>Pagurus samuelis</i> (STIMPSON) and <i>Hemigrapsus sanguineus</i> (DE HAAN).....	270
OKAZAKI, K.: Skeleton formation of sea urchin larvae. II. Organic matrix of the spicule.....	283
Hsü, C.: The effect of aureomycin on development of the frog.....	321

No. 4 (February-March, 1961)

HISHIDA, T., TOMITA, H. and YAMAMOTO, T.: Melanin forma- tion in color varieties of the medaka (<i>Oryzias latipes</i>).....	335
TOMITA, H. and HISHIDA, T.: A quantitative study on phenol oxidase of skins in color varieties of the medaka (<i>Oryzias</i> <i>latipes</i>).....	347
SAKAI, Y. T.: Method for removal of chorion and fertilization of the naked egg in <i>Oryzias latipes</i>	357
KOJIMA, M. K.: Cyclic changes of the cortex and the cyto- plasm of the fertilized and the activated sea urchin egg. III. Susceptibility of artificially activated eggs to cleavage- inducing action of hypertonic, DNP- and NaN_3 sea water.....	369
HAINO, K. and DAN, J. C.: Some quantitative aspects of the acrosomal reaction to jelly substance in the sea urchin.....	376
ÔGI, K.: Vegetalization of the presumptive ectoderm of the <i>Triturus</i> -gastrula by exposure to lithium chloride solution.....	384
AKETA, K.: Studies on the production of the fertilization acid in sea urchin eggs. I. Acid production at fertilization and activation, and the effect of some metabolic inhibitors.....	397
AKETA, K.: Studies on the production of the fertilization acid in sea urchin eggs. II. Experimental analysis of the production mechanism.....	406
TOMITA, H. and MATSUDA, N.: Deformity of vertebrae induced by lathyrogenic agents and phenylthiourea in the medaka (<i>Oryzias latipes</i>).....	413
TOMITA, H. and HISHIDA, T.: On the phenol oxidase of em- bryonic and larval stages of the medaka (<i>Oryzias latipes</i>).....	423

I-16-1,

Embryologia

Errata
for
Embryologia Vol. 5, No. 3

p. 313, in Table 4, designation of
the first column,

for Stage of transfer to 0.1 Ca read Stage of transfer to 0.7 P



THE EMBRYOLOGIA SOCIETY
Nagoya, Japan